

Logarithmic comparison theorem and $\mathcal{D}$ -modules: an overview

Tristan Torrelli¹

ABSTRACT. Let $D \subset X$ be a divisor in a complex analytic manifold. A natural problem is to determine when the de Rham complex of meromorphic forms on X with poles along D is quasi-isomorphic to its subcomplex of logarithmic forms. In this mostly expository note, we recall the main results about this problem. In particular, we point out the relevance of the theory of $\mathcal{D}$ -modules to this topic.

Introduction

Let X be a complex analytic manifold of dimension $n \geq 2$. Given a divisor $D \subset X$, we denote j the natural inclusion $X \setminus D \hookrightarrow X$. Let $\Omega_X^\bullet(\star D)$ denote the complex of meromorphic forms on X with poles along D . From the Grothendieck Comparison Theorem [17], the de Rham morphism

$$\Omega_X^\bullet(\star D) \longrightarrow \mathbf{R}j_* \mathbf{C}_{X \setminus D}$$

is a quasi-isomorphism. In particular, if $X = \mathbf{C}^n$, then for each cohomology class $c \in H^p(\mathbf{C}^n \setminus D, \mathbf{C})$, there exists a differential form $w \in \Omega_X^p(\star D)$ such that for any p -cycle σ on $\mathbf{C}^n \setminus D$, one has $c(\sigma) = \int_\sigma w$.

It is natural to ask what one can say about the form w . For example, if D is a complex submanifold then the order of the pole of w can be taken to be 1. The question of the order of the pole goes back to P.A. Griffiths [16]. We recall that a meromorphic form $w \in \Omega_X^p(\star D)$ is *logarithmic* if w and dw have at most a simple pole along D ; let $\Omega_X^\bullet(\log D) \subset \Omega_X^\bullet(\star D)$ denote the subcomplex of logarithmic forms with pole along D , introduced in full generality by K. Saito in [25]. In the initial case of normal crossing divisors, P. Deligne [14] proved that the filtered morphism $(\Omega_X^\bullet(\log D), \sigma) \hookrightarrow (\Omega_X^\bullet(\star D), P)$ where P is the pole

¹Laboratoire J.A. Dieudonné, UMR du CNRS 6621, Université de Nice Sophia-Antipolis, Parc Valrose, 06108 Nice Cedex 2, France. *E-mail:* torrelli@math.unice.fr

2000 Mathematics Subject Classification: 32C38, 32S25, 14F10, 14F40.

Keywords: de Rham complexes, $\mathcal{D}$ -modules, free divisors, hyperplane arrangements, logarithmic comparison theorem, logarithmic vector fields, Bernstein polynomial.

order filtration and σ is induced by P , is a quasi-isomorphism compatible with filtrations. This fact was crucial in order to define a mixed Hodge structure on the cohomology of a quasi-projective algebraic variety. Hence, one says that D satisfies the *logarithmic comparison theorem* if

LCT(D): The inclusion $\Omega_X^\bullet(\log D) \hookrightarrow \Omega_X^\bullet(\star D)$ is a quasi-isomorphism.

A natural problem is therefore to find classes of divisors satisfying this condition, and also to understand its meaning. Initiated by F.J. Castro-Jiménez, D. Mond and L. Narváez-Macarro [9], this problem has been intensively studied these last years. In this note, we gather together the main open questions² and the main results. Essentially, they were obtained for hypersurfaces with isolated singularities, hyperplane arrangements and free divisors (see §1). In this last case, we recall the characterization in terms of $\mathcal{D}$ -modules due to the Sevillian group around F.J. Castro-Jiménez and L. Narváez-Macarro (Theorem 2.3). Finally, we explain how enlightening this viewpoint is for the general study of the condition **LCT**(D) (see §3).

Acknowledgements. This research has been supported by a Marie Curie Fellowship of the European Community (programme FP5, contract HPMD-CT-2001-00097). The author is very grateful to the Departamento de Álgebra, Geometría y Topología (Universidad de Valladolid) for hospitality during the fellowship. Moreover, he is happy to thank Francisco Jesús Castro-Jiménez, Antoine Douai, Luis Narváez-Macarro and José María Ucha-Enríquez for judicious comments.

1 Main results about **LCT**(D)

There are few families of divisors for which this condition **LCT**(D) has been studied. Indeed, it is difficult to work directly with the complex $\Omega_X^\bullet(\log D)$ since we do not have in general a description of the logarithmic forms.

1.1 The case of weighted homogeneous hypersurfaces with an isolated singularity

We recall that a polynomial $h \in \mathbf{C}[x] = \mathbf{C}[x_1, \dots, x_n]$ is *weighted homogeneous* of weight $d \in \mathbf{Q}^+$ for a system $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbf{Q}^{*+})^n$ if h is a (nontrivial) $\mathbf{C}$ -linear combination of monomials $x_1^{\gamma_1} \cdots x_n^{\gamma_n}$ with $\sum_{i=1}^n \alpha_i \gamma_i = d$. In other words, we have the relation $\chi(h) = dh$ where χ is the Euler-vector field $\alpha_1 x_1 \partial_1 + \cdots + \alpha_n x_n \partial_n$ associated with α .

²See the appendix.

As usual, the case of weighted homogeneous polynomials defining an isolated singularity at the origin provides combinatorial formulas in terms of the weights associated with the Jacobian algebra $A_h = \mathbf{C}[x]/(h'_{x_1}, \dots, h'_{x_n})$. D. Mond and M. Holland [18] have obtained the following characterization:

THEOREM 1.1 *Assume that $n \geq 3$. Let $h \in \mathbf{C}[x]$ be a weighted homogeneous polynomial of degree d for a system $\alpha \in (\mathbf{Q}^{*+})^n$. Assume that h defines an isolated singularity at the origin. Let $D \subset \mathbf{C}^n$ be the hypersurface defined by h . The following conditions are equivalent:*

1. *The logarithmic comparison theorem holds for D .*
2. *The link of 0 in D is a $\mathbf{Q}$ -homology sphere.*
3. *There is no weighted homogeneous element in A_h whose weight belongs to the set $\{k \times d - \sum_{i=1}^n \alpha_i ; 1 \leq k \leq n - 2\} \subset \mathbf{Q}$.*

In particular, the logarithmic comparison theorem does not hold in general (see also Proposition 3.2). For example, if $h = x_1^2 + \dots + x_n^2$ then we can take $d = 2$, $\alpha_1 = \dots = \alpha_n = 1$ and $A_h = \mathbf{C}\bar{1}$. Thus $\mathbf{LCT}(D)$ is satisfied if and only if $n = 2$ or n is odd.

1.2 The case of hyperplane arrangements

Let D be a finite union of affine hyperplanes H in $X = \mathbf{C}^n$, i.e. $H = \{\alpha_H = 0\}$ where $\alpha_H \in \mathbf{C}[x_1, \dots, x_n]$ are polynomials of degree one. We can associate with D the $\mathbf{C}$ -subalgebra of $\Omega_X^\bullet(\star D)$ generated by 1 and the 1-forms $d\alpha_H/\alpha_H$. Let $R^\bullet(D)$ denote this algebra of differential forms. It is well known that $R^\bullet(D)$ is isomorphic to the so-called Orlik-Solomon algebra. Moreover

THEOREM 1.2 [2] *For all $k \geq 0$, we have $R^k(D) \cong H^k(X \setminus D, \mathbf{C})$.*

On the other hand, we can consider the following complex of $\mathbf{C}$ -vector spaces: $0 \rightarrow R^0(D) \xrightarrow{0} \dots \xrightarrow{0} R^n(D) \rightarrow 0$ as a subcomplex of $\Omega_X^\bullet(\log D)$. Thus, a natural question is: does the logarithmic comparison theorem hold for any hyperplane arrangement? This was conjectured by H. Terao in [28]. This is true for tame arrangements (such as free arrangements, generic arrangements or complex reflection arrangements) and when $n \leq 4$ (see [33]). But in general, the question is still open.

1.3 The case of free divisors

Let $\mathcal{O}_X$ be the sheaf of holomorphic functions on X . Given a divisor $D \subset X$, we will denote by $h_D \in \mathcal{O}_{X,m} \cong \mathcal{O} = \mathbf{C}\{x_1, \dots, x_n\}$ a defining equation of D at $m \in D$.

A holomorphic vector field v is *logarithmic* along D if for any point $m \in D$, $v(h_D)$ belongs to $h_D \mathcal{O}_{X,m}$. Let $\text{Der}(-\log D)$ denote the (coherent) $\mathcal{O}_X$ -module of logarithmic vector fields. We recall a property studied by K. Saito in [25].

DEFINITION 1.3 *A divisor $D \subset X$ is free at the point $m \in D$ if $\text{Der}(-\log D)_m$ is $\mathcal{O}_{X,m}$ -free. It is a free divisor if $\text{Der}(-\log D)$ is locally free.*

From the inclusions $h_D \text{Der}(\mathcal{O}_X)_m \subset \text{Der}(-\log D)_m \subset \text{Der}(\mathcal{O}_X)_m$, the rank of $\text{Der}(-\log D)$ is also equal to n .

EXAMPLE 1.4 Free divisors appear in many different contexts.

(i) Normal crossing divisors are free. Indeed, in local coordinates such that $h_D = x_1 \cdots x_p$, then

$$\text{Der}(-\log D) = \mathcal{O}_{x_1} \partial_1 \oplus \cdots \oplus \mathcal{O}_{x_p} \partial_p \oplus \mathcal{O} \partial_{p+1} \oplus \cdots \oplus \mathcal{O} \partial_n.$$

(ii) Plane curves are free (K. Saito [25]).

(iii) Complex reflection arrangements are free (H. Terao [29]). For example, the braid arrangement, defined by $\prod_{1 \leq i < j \leq n} (x_i - x_j)$ in $\mathbf{C}^n$, is free.

(iv) The discriminant of a versal deformation of an isolated complete intersection singularity is a free divisor (see [26]; [21], Corollary 6.13; [1]).

Most of the known results about the logarithmic comparison theorem have been obtained for free divisors. Indeed, the logarithmic de Rham complex $\Omega_X^\bullet(\log D)$ is also explicit. More precisely, because of the duality between $\Omega_X^1(\log D)$ and $\text{Der}(-\log D)$, $\Omega_X^1(\log D)$ is also a free $\mathcal{O}_X$ -module and we have $\Omega_X^q(\log D) = \bigwedge^q \Omega_X^1(\log D)$ for $1 \leq q \leq n$ (see [25]). Firstly, we have the following characterization for plane curves:

THEOREM 1.5 [5] *If $D \subset X = \mathbf{C}^2$ is a plane curve, then the logarithmic comparison theorem holds if and only if D is locally weighted homogeneous.*

This last condition means that for all $m \in D$, there exists an analytic change of coordinates ϕ such that $h_D \circ \phi$ is a weighted homogeneous polynomial; for example, weighted homogeneous hypersurfaces with an isolated singularity and hyperplane arrangements are locally weighted homogeneous. This unusual condition is the suitable one in this context for doing inductions on the dimension of D (see the proof of Proposition 3.1 for example). More generally, we have

THEOREM 1.6 [9] *Let $D \subset X$ be a locally weighted homogeneous free divisor. Then the logarithmic comparison theorem holds for D .*

Among the free divisors in Example 1.4, the one given in (i), (iii) and some³ of (iv) are locally weighted homogeneous.

The converse is false in general. For example, $h = x_1x_2(x_1+x_2)(x_1+x_2x_3)$ defines a free divisor $D \subset \mathbf{C}^3$ such that $\mathbf{LCT}(D)$ is true and h is not weighted homogeneous (see [5], §4). Meanwhile, h belongs to the ideal of its partial derivatives. In other words, there exists locally a vector field v such that $v(h_D) = h_D$; one says sometimes that h is *Euler-homogeneous*. In fact, we have no example of a free divisor $D = V(h)$ verifying $\mathbf{LCT}(D)$ which is not Euler-homogeneous. This is true for a Koszul-free divisor (see Definition 2.2, Theorem 3.4); moreover, M. Granger and M. Schulze [15] have obtained the following result:

THEOREM 1.7 *Let $D = V(h) \subset X = \mathbf{C}^3$ be a free divisor. If the logarithmic comparison theorem holds for D , then h is Euler-homogeneous.*

For $n \geq 4$, this question is still open (see [5], Conjecture 1.4) and it can be extended for a general divisor.

2 A differential viewpoint for free divisors

Here we recall how the condition $\mathbf{LCT}(D)$ may be interpreted in terms of $\mathcal{D}_X$ -modules for free divisors $D \subset X$, as it was initiated by F.J. Calderón-Moreno in [4].

2.1 Preliminaries

Given a complex analytic manifold X of dimension $n \geq 2$, we denote $\Omega_X^\bullet$ the complex of holomorphic differential forms on X and $(\mathcal{D}_X, F_\bullet)$ the sheaf of linear differential operators with holomorphic coefficients filtered by order. Locally at a point $m \in X$, we have $\mathcal{O}_{X,m} \cong \mathcal{O} = \mathbf{C}\{x_1, \dots, x_n\}$ and $\mathcal{D}_{X,m} \cong \mathcal{D} = \mathcal{O}\langle \partial_1, \dots, \partial_n \rangle$; moreover we identify $\mathrm{gr}^F \mathcal{D}$ with $\mathcal{O}[\xi] = \mathcal{O}[\xi_1, \dots, \xi_n]$.

The so-called Riemann-Hilbert correspondence of Z. Mebkhout and M. Kashiwara (see [20], [23], [24]) asserts that there is an equivalence of categories between the category $hr(\mathcal{D}_X)$ of (left) regular holonomic $\mathcal{D}_X$ -modules and the one of perverse sheaves $Perv_X(\mathbf{C})$ on X via the de Rham functor

$$\begin{aligned} hr(\mathcal{D}_X) &\longrightarrow Perv_X(\mathbf{C}) \\ \mathcal{M} &\longmapsto \mathrm{DR}(\mathcal{M}) = \Omega_X^\bullet \otimes_{\mathcal{O}_X} \mathcal{M}. \end{aligned}$$

³For more details, see [9].

Roughly speaking, a perverse sheaf on X is a special type of complex of sheaves on X whose cohomology groups are constructible in $\mathbf{C}$ -vector spaces of finite dimension on a stratification of X . For example, $\mathcal{O}_X$ is regular holonomic and $\mathrm{DR}(\mathcal{O}_X) = \Omega_X^\bullet$ is quasi-isomorphic to the constant sheaf $\mathbf{C}_X$ by the Poincaré lemma.

2.2 On the perversity of $\Omega_X^\bullet(\log D)$

Given a divisor $D \subset X$, we consider the sheaf $\mathcal{O}_X(\star D)$ of meromorphic functions with poles along D . As $\mathcal{O}_X(\star D)$ is regular holonomic, the meromorphic de Rham complex $\mathrm{DR}(\mathcal{O}_X(\star D)) = \Omega_X^\bullet(\star D)$ is a perverse sheaf too. Thus it is natural to investigate conditions on D in order to get the perversity of $\Omega_X^\bullet(\log D)$. In the case of free divisors, this question was studied by F.J. Calderón-Moreno and L. Narváez-Macarro in [4], [8]. They obtained the following characterization:

THEOREM 2.1 *Let $D \subset X$ be a free divisor. Then the logarithmic complex $\Omega_X^\bullet(\log D)$ is perverse if and only if the following conditions are satisfied:*

1. *the complex $\mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)}^L \mathcal{O}_X$ is concentrated in degree 0;*
2. *the $\mathcal{D}_X$ -module $\mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)} \mathcal{O}_X$ is holonomic.*

From [11], we say also that D is of *Spencer type*. Here $\mathcal{V}_0^D(\mathcal{D}_X) \subset \mathcal{D}_X$ is the coherent sheaf of rings⁴ of logarithmic operators [4] (that is, $P \in \mathcal{D}_X$ such that locally $P \cdot (h_D^k) \subset h_D^k \mathcal{O}$ for any integer k). Let us notice that this condition 1 has no clear meaning. Thus, the problem is now to find (geometrical) criteria on a free divisor to be of Spencer type (see [8], §5). The only known condition is to be a Koszul-free divisor (see [4], Theorem 4.2.1).

DEFINITION 2.2 *A free divisor $D \subset X$ is Koszul-free if there exists locally a basis⁵ $\{\delta_1, \dots, \delta_n\}$ of $\mathrm{Der}(-\log D)$ such that the sequence of principal symbols $(\sigma(\delta_1), \dots, \sigma(\delta_n))$ is $\mathrm{gr}^F \mathcal{D}$ -regular.*

For example, plane curves and locally weighted homogeneous free divisors are Koszul-free [25], [6], [7]. This notion means also that the free divisor D has a holonomic stratification in the sense of K. Saito (see [25], §3; [3], Proposition 6.3; [4], Corollary 1.9).

Finally, we make the remark that being Koszul-free is not necessary for the perversity of $\Omega_X^\bullet(\log D)$; for example, the free divisor $D \subset \mathbf{C}^3$ defined by

⁴For some results about this sheaf for a general divisor, see [27].

⁵In fact, every basis of $\mathrm{Der}(-\log D)$ satisfies this property when D is Koszul-free.

$x_1x_2(x_1+x_2)(x_1+x_2x_3)$ is not Koszul-free but $\Omega_X^\bullet(\log D)$ is perverse [4], [5]. On the other hand, the complex $\Omega_X^\bullet(\log D)$ is not perverse for any free divisor. For example, the divisor $(x_1^5 + x_2^4 + x_1^4x_2)(x_1 + x_2x_3) = 0$ is free but not of Spencer type (see [8], §5).

2.3 A differential characterization of $\mathbf{LCT}(D)$

Let us now give a differential analogue of condition $\mathbf{LCT}(D)$. F.J. Castro-Jiménez and J.M. Ucha-Enríquez began work on this problem in [10] & [11], and they obtained a characterization for free divisors of Spencer type [12]. For a general free divisor, we have the following generalization⁶ due to F.J. Calderón-Moreno and L. Narváez-Macarro [8]:

THEOREM 2.3 *Let $D \subset X$ be a free divisor. Then the inclusion*

$$\Omega_X^\bullet(\log D) \hookrightarrow \Omega_X^\bullet(\star D)$$

is a quasi-isomorphism if and only if the following conditions are satisfied:

1. *the complex $\mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)}^L \mathcal{O}_X(D)$ is concentrated in degree 0;*
2. *the natural morphism*

$$\varphi_D : \mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)} \mathcal{O}_X(D) \longrightarrow \mathcal{O}_X(\star D)$$

is an isomorphism.

Here $\mathcal{O}_X(D)$ denotes the $\mathcal{V}_0^D(\mathcal{D}_X)$ -module of meromorphic functions with at most a simple pole along D . Unfortunately, this characterization is no more explicit than condition $\mathbf{LCT}(D)$; meanwhile, condition 1 is verified by Koszul-free divisors.

A key point in the proofs of Theorems 2.3 and 2.1 is the relationship between the duals of any integral logarithmic connection over the base ring $\mathcal{D}_X$ and $\mathcal{V}_0^D(\mathcal{D}_X)$ (see [11]; [8], §3). In particular, the holonomic $\mathcal{D}_X$ -modules $\mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)} \mathcal{O}_X(D)$ and $\mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)} \mathcal{O}_X$ are dual when D is a free divisor of Spencer type, and we have $\mathrm{DR}(\mathcal{D}_X \otimes_{\mathcal{V}_0^D(\mathcal{D}_X)} \mathcal{O}_X(D)) \cong \Omega_X^\bullet(\log D)$. A generalization of this duality has been obtained in [13].

3 Towards an algebraic analogue of $\mathbf{LCT}(D)$?

In this part, we explain how useful the $\mathcal{D}_X$ -modules are in the general study of the condition $\mathbf{LCT}(D)$.

⁶In fact, they obtained this result for any integral logarithmic connection (see [8], Theorem 4.1).

3.1 Preliminaries

What can one say about the free divisor D when the morphism φ_D is an isomorphism? First, we have locally $\mathcal{V}_0^D(\mathcal{D}) = \mathcal{O}[\delta_1, \dots, \delta_n]$ where $\{\delta_1, \dots, \delta_n\}$ is a basis of $\text{Der}(-\log D)$ [4], and $\mathcal{O}(D) \cong \mathcal{V}_0^D(\mathcal{D})/\mathcal{V}_0^D(\mathcal{D})(\delta_1 + a_1, \dots, \delta_n + a_n)$ with $\delta_i(h_D) = a_i h_D$, $1 \leq i \leq n$. Hence the morphism φ_D is given locally by

$$\begin{aligned} \varphi_D : \mathcal{D}/\tilde{I}^{log} &\longrightarrow \mathcal{O}[1/h_D] \\ P + \tilde{I}^{log} &\longmapsto P \cdot \frac{1}{h_D} \end{aligned}$$

where $\tilde{I}^{log} \subset \mathcal{D}$ is the left ideal generated by $\text{Ann}_{\mathcal{D}} 1/h_D \cap F_1 \mathcal{D}$. In particular, it is bijective if and only if the two conditions

A($1/h_D$): the left ideal $\text{Ann}_{\mathcal{D}} 1/h_D$ of operators annihilating $1/h_D$ is generated by operators of order 1,

B(h_D): the $\mathcal{D}$ -module $\mathcal{O}[1/h_D]$ is generated by $1/h_D$,

are satisfied. From a well known result of M. Kashiwara (see [19], Proposition 6.2), this last condition means that -1 is the only integral root of the Bernstein polynomial of h_D . We recall that the *Bernstein polynomial* $b_h(s)$ of $h \in \mathcal{O}$ at the origin is the (nonzero) unitary polynomial $b(s) \in \mathbb{C}[s]$ of smallest degree which satisfies a functional identity of the form:

$$b(s)h^s = P(s) \cdot h^{s+1}$$

with $P(s) \in \mathcal{D}[s] = \mathcal{D} \otimes \mathbb{C}[s]$ (see [19]). This is an analytic invariant of the ideal $h\mathcal{O}$. When $h \in \mathcal{O}$ is not a unit, it is easy to check that -1 is also a root of $b_h(s)$. For example, if $h = x_1^2 + \dots + x_n^2$ then $b_h(s) = (s+1)(s+n/2)$ with the functional identity:

$$(s+1)(s+\frac{n}{2})h^s = \frac{1}{4} \left[\frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_n^2} \right] \cdot h^{s+1}.$$

3.2 The conditions **B**(h_D) and **LCT**(D)

The differential viewpoint above is relevant since it was not at all clear that for a free divisor, **LCT**(D) needs the condition **B**(h_D); in particular, every locally weighted homogeneous free divisor D satisfies the condition **B**(h_D) at any point (Theorem 1.6 with Theorem 2.3 or Proposition 3.1, or more directly [11], Theorem 5.2). Moreover, from the inclusions

$$\Omega_X^\bullet(\log D) \subset \Omega_X^\bullet \otimes (\mathcal{D} \cdot 1/h_D) \subset \Omega_X^\bullet \otimes \mathcal{O}[1/h_D] = \Omega_X^\bullet(\star D) \quad (1)$$

it appears natural that conditions **LCT**(D) and **B**(h_D) are linked for any divisor D . As an illustration, we have the following result:

PROPOSITION 3.1 *Let $D \subset X$ be a divisor which satisfies the condition $\mathbf{LCT}(D)$. Assume that one of the following conditions is satisfied:*

1. *The divisor D is free except at isolated points.*
2. *The divisor D is locally weighted homogeneous.*

Then $\mathbf{B}(h_D)$ is satisfied at any point of D .

Proof. Firstly we prove the assertion when the condition 1 is satisfied. Let $U \subset X$ be a neighborhood of a point $m \in D$ such that $D \cap U$ is free at any point different from m . Let h_D be a defining equation of D on U . From Theorem 2.3, the condition $\mathbf{B}(h_D)$ is satisfied at any point in $D \cap U - \{m\}$ - since D is free at such a point. In particular, the $\mathcal{D}_U$ -module $\mathcal{C}$ in the short exact sequence

$$0 \rightarrow \mathcal{D}_U \cdot \frac{1}{h_D} \xrightarrow{j} \mathcal{O}_U(\star D) \longrightarrow \mathcal{C} \rightarrow 0$$

is supported at m . We just have to prove that $\mathcal{C}$ is zero. The associated long exact sequence of de Rham cohomology provides the short exact sequence of $\mathbf{C}$ -vector spaces

$$0 \rightarrow H_{DR}^n(\mathcal{D}_U \cdot \frac{1}{h_D}) \longrightarrow H_{DR}^n(\mathcal{O}_U(\star D)) \longrightarrow H_{DR}^n(\mathcal{C}) \rightarrow 0.$$

In particular, $H^n(j)$ is injective. On the other hand, the condition $\mathbf{LCT}(D)$ is satisfied; hence, we deduce from the inclusions (1) of de Rham complexes that the morphisms $H^i(j) : H_{DR}^i(\mathcal{D}_U(1/h_D)) \longrightarrow H_{DR}^i(\mathcal{O}_U(\star D))$, $0 \leq i \leq n$, are surjective. Therefore $H^n(j)$ is an isomorphism, that is, $H_{DR}^n(\mathcal{C}) = 0$. From classic results about $\mathcal{D}$ -modules supported at a point (see [22] for example), $\mathcal{C}$ is necessarily zero and the condition $\mathbf{B}(h_D)$ is satisfied at any point of D .

Now, we assume that D is locally weighted homogeneous. Let us prove the assertion by induction on dimension. If $n = 2$, then D is locally defined by a (reduced) weighted homogeneous polynomial in two variables; thus we can conclude as in the proof of Proposition 3.2 below. Let us assume that $n \geq 3$ and let m denote a point in D . From [9], Proposition 2.4, there exists a neighborhood U of m such that, for each point $w \in U \cap D$, $w \neq m$, the germ of pair (X, D, w) is isomorphic to a product $(\mathbf{C}^{n-1} \times \mathbf{C}, D' \times \mathbf{C}, (0, 0))$ where D' is a locally weighted homogeneous divisor of dimension $n - 2$. Moreover, the condition $\mathbf{LCT}(D)$ implies that $\mathbf{LCT}(D')$ is satisfied on a neighborhood of the origin (see [9], Lemma 2.2). Let $h_{D'} \in \mathcal{O}_{\mathbf{C}^{n-1}, 0}$ be a local equation of D' . By using the induction hypothesis, the $\mathcal{D}_{\mathbf{C}^{n-1}, 0}$ -module $\mathcal{O}_{\mathbf{C}^{n-1}, 0}[1/h_{D'}]$ is generated by $1/h_{D'}$, thus so is $\mathcal{O}_{\mathbf{C}^{n-1} \times \mathbf{C}, 0}[1/h_{D'}]$. In particular, the condition

$\mathbf{B}(h_D)$ is satisfied at any point in $D \cap U - \{m\}$. We conclude with the first part of the proof. $\square$

In particular, $\mathbf{LCT}(D)$ implies the condition $\mathbf{B}(h_D)$ for any divisor D with isolated singularities. For a general divisor, this question is open.

3.3 The conditions $\mathbf{A}(1/h_D)$ and $\mathbf{LCT}(D)$

From [31], Proposition 1.3, the condition $\mathbf{A}(1/h)$ implies $\mathbf{B}(h)$ for any nonzero germ $h \in \mathcal{O}$; in particular, $\mathbf{A}(1/h_D)$ is a local analogue of $\mathbf{LCT}(D)$ for any Koszul-free divisor D , and more generally, for any free divisor of Spencer type (from Theorem 2.3; see also [12]). Is $\mathbf{LCT}(D)$ locally equivalent to $\mathbf{A}(1/h_D)$ in general? This question is motivated by the following significant results. Firstly, this is true for weighted homogeneous hypersurfaces with an isolated singularity.

PROPOSITION 3.2 *Let $h \in \mathbf{C}[x]$ be a weighted homogeneous polynomial. Assume that h defines an isolated singularity at the origin. Let $D \subset \mathbf{C}^n$ be the hypersurface defined by h . Then the logarithmic comparison theorem holds for D if and only if the condition $\mathbf{A}(1/h)$ is satisfied.*

Proof. Under our assumptions, the condition $\mathbf{A}(1/h)$ is in fact equivalent to $\mathbf{B}(h)$ by Theorem 3.5. On the other hand, the polynomial $b_h(s)$ is given by the formula $b_h(s) = (s+1) \prod_{q \in \Pi} (s + |\alpha| + q)$ where $\alpha \in (\mathbf{Q}^{*+})^n$ is the system of weights such that the degree of h is equal to 1, the expression $|\alpha|$ denotes the sum $\sum_{i=1}^n \alpha_i \in \mathbf{Q}^{*+}$, and $\Pi \subset \mathbf{Q}^+$ is the set of the degrees of the weighted homogeneous elements in $A_h = \mathbf{C}[x]/(h'_{x_1}, \dots, h'_{x_n})$ (see [34], §11). We recall that $n - 2|\alpha|$ is the maximal element of Π ; in particular, $\mathbf{A}(1/h)$ is satisfied if $n = 2$ and so is $\mathbf{LCT}(D)$ (Theorem 1.5). Moreover, the set Π is symmetric about $(n/2) - |\alpha|$; hence, we deduce easily that $\mathbf{B}(h)$ is equivalent to the last condition of Theorem 1.1 when $n \geq 3$. This completes the proof. $\square$

Moreover, the condition $\mathbf{B}(h_D)$ is satisfied by any hyperplane arrangement (A. Leykin [32], Theorem 5.1), and the condition $\mathbf{A}(1/h_D)$ is true for the union of a generic arrangement with a hyperbolic arrangement [31]; this agrees with Terao's conjecture (see §1.2). The general problem is still open, and condition $\mathbf{A}(1/h)$ may only be necessary. A difficulty is the lack of families of divisors which satisfy the condition $\mathbf{LCT}(D)$.

3.4 The condition $\mathbf{A}(1/h)$

Let $h \in \mathcal{O}$ be a nonzero germ such that $h(0) = 0$. We give here some results about the meaning of the condition $\mathbf{A}(1/h)$ (see [31]).

First, we have the following easy criterion:

LEMMA 3.3 *Let $h \in \mathcal{O}$ be a nonzero germ such that $h(0) = 0$. Assume that the following conditions are satisfied:*

H(h): *h belongs to the ideal of its partial derivatives;*

B(h): *-1 is the smallest integral root of $b_h(s)$;*

A(h): *the ideal $\text{Ann}_{\mathcal{D}} h^s$ is generated by operators of order 1.*

Then the ideal $\text{Ann}_{\mathcal{D}} 1/h$ is generated by operators of order 1.

Proof. By Euclidean division, we have also a decomposition

$$\text{Ann}_{\mathcal{D}[s]} h^s = \mathcal{D}[s](s - v) + \mathcal{D}[s]\text{Ann}_{\mathcal{D}} h^s$$

where v is a vector field such that $v(h) = h$ (condition **H**(h)). Moreover, under the condition **B**(h), the ideal $\text{Ann}_{\mathcal{D}} 1/h$ is obtained by fixing $s = -1$ in a system of generators of $\text{Ann}_{\mathcal{D}[s]} h^s$. $\square$

Reciprocally, what does remain true? We recall that the condition **A**($1/h$) always implies **B**(h). On the other hand, does **A**($1/h$) imply **H**(h)? This is true for isolated singularities [30], Koszul-free germs, and suspensions of unreduced plane curve $z^N + g(x_1, x_2)$ (see [31]); this question is still open. Finally, the condition **A**($1/h$) does not imply **A**(h) in general. Indeed, Calderón's example $h = x_1 x_2 (x_1 + x_2)(x_1 + x_2 x_3)$ satisfies **LCT**(D), **A**($1/h$), **B**(h), **H**(h) and not **A**(h) (see [4], [5], [6], [10], [31]). Meanwhile, condition **A**(h) is not unrealistic, since we have the following characterization of **A**($1/h$) for Koszul-free germs:

THEOREM 3.4 [31] *Let $h \in \mathcal{O}$ be a Koszul-free germ. Then the left ideal $\text{Ann}_{\mathcal{D}} 1/h$ is generated by operators of order one if and only if the conditions **H**(h), **B**(h) and **A**(h) are satisfied.*

Moreover, condition **A**(h) is satisfied when h defines an isolated singularity (see below). Thus, we have


THEOREM 3.5 [30] *Let $h \in \mathcal{O}$ be a germ of holomorphic function defining an isolated singularity. Then the ideal $\text{Ann}_{\mathcal{D}} 1/h$ is generated by operators of order one if and only if the germ h is weighted homogeneous and the condition **B**(h) is satisfied.*

In fact, the condition **A**(h) may be considered almost as a geometric condition. Indeed, the following condition implies **A**(h):

$\mathbf{W}(h)$: the relative conormal space W_h is defined by linear equations in ξ since $W_h = \overline{\{(x, \lambda dh) : \lambda \in \mathbf{C}\}} \subset T^*\mathbf{C}^n$ is the characteristic variety of $\mathcal{D}h^s$ [19]. For example, $\mathbf{W}(h)$ is true for hypersurfaces with an isolated singularity [34] and for locally weighted homogeneous free divisors [6].

Appendix

The following diagram summarizes the expected relations between the conditions studied in this note:



Here D denotes a general divisor, and the solid headed arrows represent the open questions.

References

- [1] ALEKSANDROV A. G., *Euler-homogeneous singularities and logarithmic differential forms*, Ann. Global Anal. Geom. 4 (1986) 225–242.
- [2] BRIESKORN E., *Sur les groupes de tresses [d’après V. I. Arnold]*, séminaire Bourbaki, 24ème année (1971/1972), Exp. No. 401, pp. 21–44. Lect. Notes in Math., Vol. 317, Springer, Berlin, 1973.
- [3] BRUCE F.W., ROBERTS R.M., *Critical points of functions on analytic varieties*, Topology 27 (1988) 57–90.
- [4] CALDERÓN-MORENO F.J., *Logarithmic differential operators and logarithmic de Rham complexes relative to a free divisor*, Ann. Sci. École Norm. Sup. 32 (1999) 577–595.

- [5] CALDERÓN-MORENO F.J., CASTRO-JIMÉNEZ F.J., MOND D., NARVÁEZ-MACARRO L., *Logarithmic cohomology of the complement of a plane curve*, Comment. Math. Helv. 77 (2002) 24–38.
- [6] CALDERÓN-MORENO F.J., NARVÁEZ-MACARRO L., *The module $\mathcal{D}f^s$ for locally quasi-homogeneous free divisors*, Compos. Math. 134 (2002) 59–74.
- [7] CALDERÓN-MORENO F.J., NARVÁEZ-MACARRO, L., *Locally quasi-homogeneous free divisors are Koszul free*, Tr. Mat. Inst. Steklova 238 (2002) 81–85
- [8] CALDERÓN-MORENO F.J., NARVÁEZ-MACARRO L., *Dualité et comparaison sur les complexes de de Rham logarithmiques par rapport aux diviseurs libres*, Ann. Inst. Fourier 55 (2005) 47–75.
- [9] CASTRO-JIMÉNEZ F.J., MOND D., NARVÁEZ-MACARRO L., *Cohomology of the complement of a free divisor*, Trans. Amer. Math. Soc. 348 (1996) 3037–3049.
- [10] CASTRO-JIMÉNEZ F.J., UCHA-ENRÍQUEZ J.M., *Explicit comparison theorems for $\mathcal{D}$ -modules*, J. Symbolic Comput. 32 (2001) 677–685.
- [11] CASTRO-JIMÉNEZ F.J., UCHA-ENRÍQUEZ J.M., *Free divisors and duality for $\mathcal{D}$ -modules*, Tr. Mat. Inst. Steklova 238 (2002) 97–105.
- [12] CASTRO-JIMÉNEZ F.J., UCHA-ENRÍQUEZ J.M., *Testing the logarithmic comparison theorem for free divisors*, Experiment. Math. 13 (2004) 441–449.
- [13] CASTRO-JIMÉNEZ F.J., UCHA-ENRÍQUEZ J.M., *Quasi-free divisors and duality*, C. R. Acad. Sci. Paris 338 (2004) 461–466.
- [14] DELIGNE P., *Équations différentielles à points singuliers réguliers*, Lect. Notes in Math. 163 (1970).
- [15] GRANGER M., SCHULZE M., *On the formal structure of logarithmic vector fields*, arXiv.org math.AG/0412014. To appear in Compos. Math.
- [16] GRIFFITHS P.A., *On the periods of certain rational integrals I*, Ann. of Math. 90 (1969) 460–495.
- [17] GROTHENDIECK A., *On the de Rham cohomology of algebraic varieties*, Pub. Math. I.H.E.S. 29 (1966) 95–105.
- [18] HOLLAND M., MOND D., *Logarithmic differential forms and the cohomology of the complement of a divisor*, Math. Scand. 83 (1998) 235–254.

- [19] KASHIWARA M., *B-functions and holonomic systems*, Invent. Math. 38 (1976) 33–53.
- [20] KASHIWARA M., *The Riemann-Hilbert problem for holonomic systems*, Publ. RIMS, Kyoto Univ. 20 (1984) 319–365.
- [21] LOOIJENGA E.J.N., *Isolated singular points on complete intersections*, London Mathematical Society, Lecture Note Series 77 (1984).
- [22] MALGRANGE B., *Le polynôme de Bernstein d’une singularité isolée*, Lect. Notes in Math. 459 (1975) 98–119.
- [23] MEBKHOUT Z., *Une équivalence de catégories. Une autre équivalence de catégories*, Compos. Math. 51 (1984) 51–88.
- [24] MEBKHOUT Z., *Le théorème de positivité, le théorème de comparaison et le théorème d’existence de Riemann*, Séminaires et Congrès 8, Soc. Math. France, Paris (2004) 165–310.
- [25] SAITO K., *Theory of logarithmic differential forms and logarithmic vector fields*, J. Fac. Sci. Univ. Tokyo 27 (1980) 265–291.
- [26] SAITO K., *Primitive forms for a universal unfolding of a function with an isolated critical point*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 28 (1982) 775–792.
- [27] SCHULZE M., *A criterion for the logarithmic differential operators to be generated by vector fields*, arXiv.org math.CV/0406023.
- [28] TERAOKA H., *Forms with logarithmic pole and the filtration by the order of the pole*, Proceedings of the International Symposium on Algebraic Geometry (Kyoto University, Kyoto, 1977) (Kinokuniya Book Store, Tokyo, 1978) 673–685.
- [29] TERAOKA H., *Free arrangements of hyperplanes and unitary reflection groups*, Proc. Japan Acad. Ser. A Math. Sci. 56 (1980) 389–392.
- [30] TORRELLI T., *Polynômes de Bernstein associés à une fonction sur une intersection complète à singularité isolée*, Ann. Inst. Fourier 52 (2002) 221–244.
- [31] TORRELLI T., *On meromorphic functions defined by a differential system of order 1*, Bull. Soc. Math. France 132 (2004) 591–612.
- [32] WALTHER U., *Bernstein-Sato polynomial versus cohomology of the Milnor fiber for generic hyperplane arrangements*, Compos. Math. 141 (2005) 121–145.

- [33] WIENS J., YUZWINSKY S., *De Rham cohomology of logarithmic forms on arrangements of hyperplanes*, Trans. Amer. Math. Soc. 349 (1997) 1653–1662.
- [34] YANO T., *On the theory of b-functions*, Publ. R.I.M.S. Kyoto Univ. 14 (1978) 111–202.